

STAGING

I < 2cm + NO LYMPH

II > 2cm + NO LYMPH

III SPREAD TO UTERINE VAG OR UNILAT NODES INVOLVED

IV SPREAD PERINEAL AREA AND/OR DISTAL NODES INVOLVED

MASS DISCHARGE BLEEDING PRURITIS

COMMONLY 95% SQUAMOUS

STAGING & PAINFUL D VULVA IMMUNOSUP

GI TRACT ENDOMETRIAL CERVIX

USUALLY 20 CM FROM

BLEEDING DISCHARGE MASS/ULCER

VULVAL ITCHING SORENESS IFTNES THICKENED HYPO/HYPER PIGMENTED

VULVAL & VAGINAL DISORDERS

BARTHOLIN'S GLAND (MUCOUS BLEEDING CERVIX M M M M M)

TO EXCISE BIOPSY TOPICAL STEROID

SUPERFICIAL

SIMPLEX LICHEN PLANUS

MANAGEMENT AVOID IRRITANTS -> SOAPS EMOLIENTS TOPICAL STEROID ANTIH1S

3P'S SMALL CURRISH PAPULAR LES + PLAIN TOP

ANOGENITAL AREA

VULVAL EPITHELIUM PRURITIS SORENESS

INFECTED CAN BE PAINFUL ABS BLEBBERED CYST

ACUTELY PAINFUL LARGE, RED TENDER SWELLING



